

RIVERSIDE SCHOOL



Infection Control Policy

APPROVED BY GOVERNORS

RESPONSIBLE PERSON – HEADTEACHER

Contents

Statement of Intent

1. Legal Framework

Preventative Measures

2. Ensuring a Clean Environment

3. Pupil Immunisation

4. Staff Immunisation

5. Contact with Pets and Animals

6. Water Based Activities

In the Event of Infection

7. Preventing the Spread of Infection

8. Vulnerable Pupils

9. Procedures for Unwell Pupils and Staff

10. Exclusion

11. Medication

12. Outbreaks of Infectious Diseases

13. Pregnant Staff Members

14. Staff Handling Food

15. Managing Specific Infectious Diseases

16. Monitoring and Review

Appendix 1 – Managing Specific Infectious Diseases

Appendix 2 – Infection Absence Periods

Appendix 3 – Diarrhoea and Vomiting Outbreak Action Checklist

Statement of Intent

Infections in schools can spread easily in a school due to:

- Pupils' undeveloped immune systems;
- The close-contact nature of the environment;
- Some pupils having not yet received full vaccinations; and
- Pupils' poor understanding of good hygiene practices.

Infections commonly spread in the following ways:

- **Respiratory spread** – contact with coughs or other secretions from an infected person.
- **Direct contact spread** – direct contact with the infecting organism e.g., skin-on-skin contact during sports.
- **Gastrointestinal spread** – contact with contaminated food or water, or contact with infected faeces or unwashed hands.
- **Blood-borne virus spread** – contact with infected blood or bodily fluids e.g., via bites or used needles.

The school actively prevents the spread of infection via the following measures:

- Maintaining high standards of personal hygiene and practice;
- Maintaining a clean environment;
- Routine immunisations; and
- Taking appropriate action when infection occurs.

This policy aims to help school staff prevent and manage infections in school. IT is not intended to be used as a tool for diagnosing disease, but rather a series of procedures informing staff what steps to take to prevent infection, and what actions to take when infection occurs.

1. Legal Framework

This policy has due regard to legislation including, but not limited to the following:

- The Control of Substances Hazardous to Health Regulations (COSHH) 2002 (amended 2004);
- Health and Safety at Work etc. Act 1974;
- The Management of Health and Safety at Work Regulations 1999;
- The Reporting of Injuries, Diseases and Dangerous Occurrences Regulations (RIDDOR) 2013;
- The Health Protection (Notification) Regulations 2010; and
- The Health Protection (Local Authority Powers) Regulations 2010.

This policy has due regard to statutory guidance including, but not limited to, the following:

- UK Health and Security Agency (2023) 'Health protection in children and young people settings, including education'; and
- DfE (2015) 'Supporting pupils at school with medical conditions'.

This policy operates in conjunction with the following school policies and documents:

- Health and Safety Policy;
- Supporting Pupils with Medical Needs in School Policy - Administration of Medication and Enteral Feeds;
- Animals in School Policy;
- First Aid Policy; and
- Ill Health and Infectious Disease Risk Assessment.

Preventative Measures

2. Ensuring a Clean Environment

Handwashing

The school will ensure all staff and pupils have access to liquid soap, warm water and paper towels. Staff will check, encourage and supervise handwashing where appropriate.

All staff and pupils will be advised to wash their hands after using the toilet, before eating or handling food, after playtime and after touching animals.

Cleaning

All cleaning staff will be appropriately trained and appropriate PPE e.g., gloves, aprons and surgical masks, will be available. The Facilities Manager will devise a cleaning schedule that clearly describes the activities required, the frequency of cleaning and who

will carry out which activities. Cleaning standards will be regularly monitored to ensure effectiveness and that all areas or surfaces in contact with food, dirt or bodily fluids are regularly cleaned and disinfected.

A cleaning contractor may be employed to carry out or support the rigorous cleaning of the premises. Cleaning equipment will be maintained to a high standard and is colour coded according to area of use. The Facilities Manager will be responsible for monitoring cleaning standards and discussing any issues that may arise with the Cleaners/ Contractors.

Sanitary Facilities

Wall mounted soap dispensers will be used in all toilets – bar soap is never used.

A foot operated waste-paper bin will be available where disposable paper towels are used. Alternatively, lid-less bins will be provided where practical.

Toilet paper will always be available in cubicles. Suitable sanitary disposal facilities will be provided where necessary, including where there are female staff and pupils aged nine and above.

Nappy Changing Areas/ Hygiene Rooms

A designated changing area will be established away from play facilities and food and drink areas, and with appropriate hand washing facilities.

Children's skin will be cleaned with disposable wipes, and nappy creams and lotions will be labelled with the relevant child's name and not shared with others.

Changing tables will be wiped with antibacterial wipes or a mild detergent wipe after each use.

Handwashing facilities are available in the room and soiled pads are disposed of inside a wrapped plastic bag and placed in the hygiene disposal bin. PPE (gloves and aprons) are provided for use during personal intimate care.

Laundry

Should any laundry need to be washed, it will be washed using the dedicated machines available.

Manual sluicing of clothing will not be permitted, and gloves and aprons will be worn when handling soiled linen or clothing. Hands will be thoroughly washed after gloves are removed.

Ventilation

Indoor spaces will be kept well ventilated to help reduce the amount of respiratory germs. Areas of the school where there may be poor ventilation will be identified e.g., through the use of CO2 monitors, and appropriate action taken, e.g., partially opening windows and doors to let fresh air in. The need for increased ventilation will always be balanced against the need to maintain a comfortable temperature for staff, pupils and visitors.

Toys and Equipment

When purchasing toys, the school will ensure they all carry a BS, BSI or CE mark and that, where possible, they can be easily cleaned. Toys will be stored in clean containers. Pupils will not be allowed to take them into toilet areas.

Regular cleaning of toys and equipment is in place. Toys that are “soft” e.g., modelling clay, will be discarded whenever they look dirty and are single use/ pupil restricted during any serious outbreak.

Water play troughs will be emptied, washed with detergent and hot water, dried and stored upside-down when not in use for long periods. When in use, the water will be replenished, at a minimum, on a daily basis, and the trough will remain covered overnight.

Managing Cuts, Bites and Bodily Fluid Spills

Standard precautions will always be taken when dealing with any cuts and abrasions.

Any spillages of blood, faeces, saliva, vomit or nasal discharges will be cleaned immediately in line with the Bodily Fluids Risk Assessment – the school spillage kit is stored in the Facilities Manager’s office.

PPE will be worn where there is a risk of contamination with blood or bodily fluids during an activity. Gloves are disposable, non-powdered vinyl or latex and CE Marked. If there is a risk of splashing to the face, disposable eye protection or reusable eye protection that is decontaminated prior to next use, will be worn.

Cuts and abrasions will be cleaned under running water or using a disposable container with water and wipes. The wound will be carefully dabbed dry then covered with a waterproof dressing or plaster. The dressing will be changed as often as is necessary. Staff will wear disposable gloves when in contact with any accident or injury e.g., washing grazes, or dressing wounds.

If a pupil suffers a bite or scratch that does not break the skin, the affected area will be cleaned with soap and warm running water. If a bite, scratch or puncture injury breaks the skin or may have introduced someone else’s blood, the affected area will be washed thoroughly with soap and warm running water, the incident will be recorded in the pupil

accident log, the wound will be covered with a waterproof dressing, and medical advice sought immediately.

When coughing or sneezing, all staff and pupils will be encouraged to cover their nose and mouth with a disposable tissue and dispose of the tissue after use, and to wash their hands afterwards.

Safe Management of Waste – Including Sharps

The school will ensure that all waste produced is dealt with by a licensed waste management company.

Any PPE used will be placed in a refuse bag and disposed of as normal domestic waste. PPE should not be put in a recycling bin or dropped as litter.

Injuries incurred through sharps found on school grounds will be treated in line with the school's Health and Safety Policy. All sharps found on school premises will be disposed of in the sharps bin whilst wearing PPE.

The School Business Manager must be informed if needles are found on site that have not been disposed of appropriately.

3. Pupil Immunisation

The school is aware of the vital role it plays in supporting the routine immunisation programme and will liaise with local health and services to share information with parents at key points.

Each pupil's immunisation status will be checked upon school entry and at the time of any in-school vaccination.

The school will support school-based immunisation programmes by hosting school nurses and School Age Immunisation Services (SAIS) and helping them with various aspects of the vaccination process, including:

- Providing space and time in the timetable for vaccination;
- Reminding staff and pupils about the date of the immunisation session(s);
- Sharing information leaflets and consent forms with pupils; and
- Providing a list of eligible pupils and their parent contact details to the SAIS team.

The school will keep up-to-date with national and local immunisation scheduling and advice via www.nhs.uk/conditions/vaccinations/.

The school will work with local health professionals to signpost parents to where they can book coronavirus (COVID-19) vaccination appointments for pupils online, at a vaccination centre or pharmacy, or at a local walk-in centre without an appointment.

The school will work with local health services and SAIS teams to ensure the effective planning and delivery of the immunisation service.

The school will permit time off for pupils to receive immunisations, where necessary.

The school will notify its regional DfE team of any anti-vaccination activity e.g., campaign letters and emails spreading misinformation about vaccination programmes. Only information from trusted sources e.g., the NHS, and where its authenticity is assured will be shared by the school.

Some vaccinations may involve an exclusion period in which pupils are not required to attend school. The administering healthcare team will provide advice in such cases.

The school will ensure that any pupils with existing medical conditions are medically cleared to receive the vaccine in question.

Whilst the school will encourage parents to have their child immunised, parental consent will always be sought before a vaccination is given.

4. Staff Immunisation

All staff will complete a medical questionnaire prior to employment, which confirms they are up-to-date with their immunisations. Staff will be encouraged to check their immunisation records and contact their GP practice if they are unsure if they are up-to-date or if they need to catch up.

Staff should be up-to-date with their immunisations; in particular we encourage the following:

- **MMR vaccination** – The MMR vaccine is the safest and most effective way to protect against measles, mumps and rubella, which are viral infections that can quickly spread and cause outbreaks. The school will liaise with staff to ensure they have had two doses of the MMR Vaccine. Female staff of childbearing age will be encouraged to check their immunisation status and contact their GP for any further medical advice.
- **Hepatitis B** – The Hepatitis B vaccine is recommended for staff who are involved in the care of individuals with severe learning disability or challenging behaviour who live in institutional accommodation. In all circumstances the school will undertake an occupational health risk assessment and pay for the vaccine if required.

Staff will be encouraged to have any immunisations in the school holidays.

5. Contact with Pets and Animals

Animals in schools will be strictly controlled and risk assessments are completed before any animal visits.

The Headteacher will assign a member of staff with suitable knowledge and experience to be responsible for animals and abide by the Animal Welfare Act 2006, which places a duty on animal owners to ensure their animal's welfare needs are met.

Animals in school will only be permitted in the following areas:

- Classrooms;
- The playground; and
- The main hall.

Animals will always be supervised when in contact with pupils and anyone handling animals will wash their hands immediately after touching them, their bedding or equipment.

Pregnant staff will be advised to avoid contact with any animal litter trays on the school premises due to the risk of toxoplasmosis.

Only organised providers or pets as therapy animals will be allowed in school and risk assessments from the organisation will be reviewed. Animals are always supervised when in contact with children and anyone handling animals will wash their hands immediately afterwards.

Visits to farms and zoos will be suitably risk assessed and are controlled by the policies and protocols covering learning outside the classroom and by risk assessment.

6. Water-Based Activities

Hydrotherapy

Hydrotherapy is governed by the control measures outlined in our Hydrotherapy Risk Assessment.

Pupils who have experienced vomiting or diarrhoea are not to use the hydrotherapy pool or attend swimming lessons for a full week after recovery.

Other Activities

Alternative water-based activities will only be undertaken at reputable centres.

Pupils and staff will cover all cuts, scratches and abrasions with waterproof dressings before taking part and will wash their hands immediately after the activity. No food or

drink will be consumed until hands have been washed.

After canoeing or rowing, staff and pupils will immediately wash or shower.

If a member of staff or a pupil becomes ill within three to four weeks of an activity taking place, the school will encourage them or their parents to seek medical advice and inform the treating doctor of their child's participation in these activities.

In the Event of Infection

7. Preventing the Spread of Infection

Parents will not bring their child to school in the following circumstances:

- The child shows signs of being unwell and needing one-to-one care;
- The child has taken, or needs to take, infant paracetamol, ibuprofen or 'Calpol';
- The child has a high temperature or fever;
- The child has been vomiting and/ or had diarrhoea within the last 48 hours; or
- The child has an infection and the recommended exclusion period, stated in Section 15 – Managing Specific Infectious Diseases, has not yet passed.

8. Vulnerable Pupils

Pupils with impaired immune defence mechanisms, known as immunosuppressed, are more likely to acquire infections. In addition, the effect of an infection is likely to be more significant for such pupils. These pupils may have a disease that compromises their immune system or be undergoing treatment e.g., chemotherapy, that has a similar effect.

The school nurse will be notified if a pupil is "vulnerable". Parents are responsible for notifying the school if their child is classified as vulnerable.

If a vulnerable pupil is thought to have been exposed to an infectious disease, the pupil's parents will be informed and encouraged to seek medical advice from their doctor or specialist.

9. Procedures for Unwell Pupils and Staff

Staff will be required to know the warning signs of pupils becoming unwell including, but not limited to, the following:

- Not being themselves;
- Not eating e.g., at break and lunch times;
- Wanting more attention or sleep than usual; and
- Displaying physical signs of being unwell e.g., watery eyes, a flushed face or clammy skin.

Where a staff member identifies a pupil as being unwell, the pupil will be taken to the first aider or the nurse's office, where their temperature will be taken by the school nurse, and the pupil's parents informed of the situation.

Where the school nurse is unavailable, staff will:

- Attempt to cool the pupil down if they are too hot, by opening a window and suggesting that the pupil removes their top layers of clothing;
- Provide the pupil with a drink of water;
- Move the pupil to a quieter area of the classroom or school;
- Ensure there is a staff member available to comfort the pupil; and
- Summon emergency medical help if required.

Pupils and staff displaying any of the signs of becoming unwell outlined above will be sent home, and the school will recommend that they see a doctor.

If a pupil is identified with sickness and diarrhoea, the pupil's parents will be contacted immediately and the child will be sent home, and may only return after 48 hours have passed without symptoms.

If a staff member is suffering from vomiting and diarrhoea, they will be sent home and may not return until 48 hours have passed without symptoms.

If the school is unable to contact a pupil's parents in any situation, the pupil's alternative emergency contacts will be contacted.

Contaminated clothing

If the clothing of the first aider or a pupil becomes contaminated, the clothing will be removed as soon as possible and placed in a plastic bag. The pupil's clothing will be sent home with the pupil, and parents are advised of the best way to launder the clothing.

Contaminated clothing will be washed separately in a washing machine, using a pre-wash cycle on the hottest temperature that the clothes will tolerate.

10. Exclusion

Pupils and staff who are showing the symptoms of an infectious disease or have been diagnosed by a health professional or diagnostic test will be advised to stay away from the school for the minimum period recommended, if required, and until well enough to return.

The school will expect parents to agree that, if their child is unwell and has symptoms of an infectious illness, such as a fever, they should not attend the school, given the potential risk to others.

If a parent insists on a pupil with symptoms attending the setting, where they have a confirmed or suspected case of an infectious illness, the school will take the decision to exclude the pupil from the school – on medical grounds – if, in the school's reasonable judgement, it is necessary to protect other pupils and staff from possible infection.

For some infections, individuals may be advised to remain away from school for a longer period of time and the school will follow any advice received from the local health protection team (HPT).

If a pupil or member of staff is a close contact of someone unwell with an infectious disease, but is not confirmed to be infected, this is not normally a valid reason for exclusion; however, the local HPT may advise on specific precautions to take in response to a case or outbreak.

The school will seek to provide remote education where a pupil is well enough to participate in it.

The school is aware that exclusion may cause challenges for parents due to unexpected time off and, that some children may become vulnerable to domestic abuse or neglect during times where they would usually be at school. When recommending exclusion on public health grounds, the school will work with their HPT to consider any adverse effects or hidden harms a pupil may be exposed to by imposing isolation, and staff will be alert and proactive in sharing information as early as possible.

The school is aware that exclusion on public health grounds may cause some pupils or staff members to feel isolated or anxious. In such situations, the school will signpost them to mental health and wellbeing support services.

11. Medication

Where a pupil has been prescribed medication by a doctor, dentist, nurse or pharmacist, the first dose will be given at home, in case the pupil has an adverse reaction.

The pupil will only be allowed to return to school 24 hours after the first dose of medication, to allow it time to take effect.

All medicine provided in school will be administered in line with the Supporting Pupils with Medical Needs in School Policy - Administration of Medication and Enteral Feeds.

12. Outbreaks of Infectious Diseases

An incident is classed as an 'outbreak' where two or more people experiencing a similar illness are linked in time or place, or a greater than expected rate of infection is present compared with the usual background rate, for example:

- Two or more pupils in the same classroom are suffering from vomiting and

diarrhoea;

- A greater number of pupils than usual are diagnosed with scarlet fever; or
- There are two or more cases of measles at the school.

Where an outbreak is suspected (even if it cannot be confirmed), the Headteacher will promptly contact the HPT to discuss the situation and agree if any actions are needed. The school will support the HPT's identified control measures with clear and prompt communication with parents and rapid coordination or arrangements e.g., staff immunisation.

The Headteacher will provide the following information:

- The number of staff and children affected;
- The symptoms present;
- The date symptoms first appeared; and
- The number of classes affected.

The local HPT will provide the school with draft letters and factsheets to distribute to parents.

The HPT will always treat outbreaks in the strictest confidence therefore, information provided to parents during an outbreak will never include names and other personal details.

If a member of staff suspects the presence of an infectious disease in the school, they will contact the school nurse for further advice.

If a parent informs the school that their child carries an infectious disease, other pupils will be observed for similar symptoms by their teachers and the school nurse.

A pupil returning to the school following an infectious disease will be asked to contact the school nurse.

If a pupil is identified as having a notifiable disease, as outlined in Appendix 2 (Infection Absence Periods), the school will inform the parents, who should inform their child's GP. It is a statutory requirement for doctors to then notify their local UK Health Security Agency Centre.

During an outbreak, enhanced or more frequent cleaning protocols may be undertaken, in line with advice provided by the local HPT. The School Business Manager will liaise with the cleaning contractor to ensure these take place.

Under the Health Protection Regulations 2010, the school will always report instances of the following diseases to the HPT:

- Acute encephalitis;

- Acute meningitis;
- Acute poliomyelitis;
- Acute infectious hepatitis;
- Anthrax;
- Botulism;
- Brucellosis;
- Cholera;
- Coronavirus (COVID-19);
- Diphtheria;
- Enteric fever (typhoid or paratyphoid fever);
- Food poisoning;
- Haemolytic uraemic syndrome (HUS);
- Infectious bloody diarrhoea;
- Invasive group A streptococcal disease and scarlet fever;
- Legionnaires' disease;
- Leprosy;
- Malaria;
- Measles;
- Meningococcal septicaemia;
- Mumps;
- Plague;
- Rabies;
- Rubella;
- SARS;
- Smallpox;
- Tetanus;
- Tuberculosis;
- Typhus;
- Viral haemorrhagic fever (VHF)
- Whooping cough; and
- Yellow fever.

13. Pregnant Staff Members

If a pregnant staff member develops a rash, or is in direct contact with someone who has a potentially contagious rash, the school will strongly encourage them to speak to their GP or midwife.

Pregnant staff members will be advised to ensure they are up-to-date with the recommended vaccinations, including those against coronavirus.

Chickenpox

If a pregnant staff member has not already had chickenpox or shingles, becoming infected can affect the pregnancy. If a pregnant staff member believes they have been exposed to chickenpox or shingles and have not had either infection previously, they will

be advised to speak to their midwife or GP as soon as possible. If a pregnant staff member is unsure whether they are immune, the school will encourage them to take a blood test.

Measles

If a pregnant staff member is exposed to measles, they will inform their midwife immediately. All female staff under the age of 25, who work with young children, are asked to provide evidence of two doses of MMR vaccine or a positive history of measles.

Rubella (German Measles)

If a pregnant staff member is exposed to rubella, they will inform their midwife immediately. All female staff under the age of 25, who work with young children, are asked to provide evidence of two doses of MMR vaccine or a positive history of rubella.

Slapped Cheek Disease (Parvovirus B19)

If a pregnant staff member is exposed to slapped cheek disease they will inform their midwife promptly.

14. Staff Handling Food

Food handling staff suffering from transmittable diseases will be excluded from all food handling activity until advised by the local Environmental Health Officer (EHO) that they are clear to return to work. Both food handling staff and midday assistants are not permitted to attend work if they are suffering from diarrhoea and/ or vomiting. They are not permitted to return to work until 48 hours have passed since diarrhoea and/ or vomiting occurred, or until advised by the local EHO that they are allowed to return to work.

The school will notify the local Environmental Health Department as soon as we are notified that a staff member engaged in the handling of food has become aware that they are suffering from, or likely to be carrying, an infection that may cause food poisoning.

Food handlers are required by law to inform the school if they are suffering from any of the following:

- Typhoid fever;
- Paratyphoid fever;
- Other salmonella infections;
- Dysentery;
- Shigellosis;
- Diarrhoea (where the cause of which has not been established);
- Infective jaundice;
- Staphylococcal infections likely to cause food poisoning like impetigo, septic skin lesions, exposed infected wounds, boils; and

- E.coli VTEC infection.

Formal exclusions will be issued where necessary, but employees are expected to provide voluntary 'off work' certificates from their GP.

15. Managing Specific Infectious Diseases

When an infectious disease occurs in the school, staff will follow the appropriate procedures set out in Appendix 1 – Managing Specific Infectious Diseases.

16. Monitoring and Review

All members of staff will be required to familiarise themselves with this policy as part of their induction programme.

The Headteacher will review this policy on an annual basis and will make any changes necessary, taking into account the current effectiveness of infection control and prevention.

Appendix 1 – Managing Specific Infectious Diseases

Disease	Symptoms	Considerations	Exclusion Period
Athlete's Foot	Scaling, peeling or cracking of the skin, particularly between the toes and on soles of the feet, or blisters containing fluid. The infection may be itchy, and toenails can become discoloured, thick or crumbly.	Cases are advised to see their local pharmacy or GP for advice and treatment	Exclusion is not necessary.
Chicken Pox	Sudden onset of fever with a runny nose, cough and generalised rash. The rash then blisters and scabs over. Several blisters may develop at once, so there may be scabs in various stages of development. Blisters typically crust up and fall off naturally within one to two weeks. Some mild infections may not present symptoms.	Cases are advised to consider pharmacy remedies to alleviate symptoms and consult their GP. Immediate medical advice should be sought if abnormal symptoms develop, e.g., infected blisters, chest pain or difficulty breathing.	Chickenpox is infectious from 48 hours prior to a rash appearing, and until all blisters have crusted over, typically five to six days after the onset of a rash. Cases will be excluded from school for at least five days from the onset of a rash and until all blisters have dried and crusted over. It is not necessary for all the spots to have healed before the case returns to school.
Cold Sores	The first signs of cold sores are tingling, burning or itching in the affected area. Around 24 hours after the first signs appear the area will redden and swell, resulting in a fluid-filled blister or blisters. After blistering, they may form ulcers, then dry up and crust over.	Cases are advised not to touch the cold sore, or pick at the blisters. Sufferers of cold sores should avoid kissing people and should not share food and items such as cutlery, cups, towels and facecloths.	Exclusion is not necessary.
Conjunctivitis	The eye(s) become reddened and swollen, and there may be a sticky or watery discharge. Eyes may feel itchy	Cases are encouraged to seek advice, wash their hands frequently and not to rub their eyes.	Exclusion is not necessary. In the case of an unmanageable

Disease	Symptoms	Considerations	Exclusion Period
	and 'gritty'.	Parents will be advised to seek advice and treatment from their local pharmacist.	outbreak, exclusion may become necessary, as per the HPT's advice.
Cryptosporidiosis	Symptoms include abdominal pain, diarrhoea and occasionally vomiting.	Staff and pupils will be asked to wash hands frequently. Kitchen and toilet areas will be cleaned regularly.	Cases will be excluded until 48 hours have passed since symptoms were present.
Diarrhoea and vomiting (Gastroenteritis)	Symptoms include diarrhoea and/ or vomiting; diarrhoea is defined as three or more liquid or semi-liquid stools in a 24-hour period.	The HPT will be contacted where there are more cases than usual.	Cases will be excluded until 48 hours have passed since symptoms were present – for some infections, longer periods are required, the HPT will advise accordingly. If medication is prescribed, the full course must be completed and there must be no further symptoms displayed for 48 hours following completion of the course before the cases may return to school. Cases will be excluded from swimming for two weeks following their last episode of diarrhoea.
E. coli STEC	Symptoms vary but include diarrhoea which can be bloody, abdominal pain, vomiting and fever.	Cases will immediately be sent home and advised to speak to their GP. The HPT will be consulted in all cases.	Cases will be excluded whilst symptomatic and for 48 hours after symptoms have resolved. Where the sufferer poses an increased risk e.g., food handlers, pre-school infants, they will be excluded until a negative stool sample has been confirmed.
Food Poisoning	Symptoms normally appear within one	Cases will be sent home.	

Disease	Symptoms	Considerations	Exclusion Period
	to two days of contaminated food being consumed although they may start at any point between a few hours and several weeks later. The main symptoms are likely to be nausea, vomiting, diarrhoea, abdominal pain and fever.	The HPT will be contacted where two or more cases with similar symptoms are reported. All outbreaks of food poisoning outbreak will be investigated.	Cases will be excluded until 48 hours have passed since symptoms were present. For some infections, longer exclusion periods may be required. The HPT will advise in such cases.
Giardiasis	Infection can be asymptomatic and the incubation period is between 5 and 25 days. Symptoms can include abdominal pain, bloating, fatigue and pale, loose stools.	Cases will be sent home. The HPT will be contacted where two or more cases with similar symptoms are reported.	Cases will be excluded until 48 hours have passed since symptoms were present.
Glandular Fever	Symptoms include severe tiredness, aching muscles, sore throat, high fever, swollen glands in the neck and occasionally jaundice.	The sufferer may feel unwell for several months with fatigue and the school will provide reasonable adjustments where necessary.	Exclusion is not necessary, and cases can return to school as soon as they feel well.
Group A Streptococcus (GAS)	Symptoms include flu-like symptoms, sore throat, rough rash, scabs and sores (impetigo), pain and swelling, severe muscle aches, nausea and vomiting.	GAS can cause a number of infections, some mild and some more serious. Milder infections can be easily treated with antibiotics and usually recover at home in a few days.	Cases will be excluded for 24 hours after starting to take antibiotics.
Hand, Foot and Mouth Disease	Symptoms include a fever, reduced appetite and generally feeling unwell. One or two days later, a rash with blisters may develop with blisters on the inside of cheeks, gums, sides of the tongue, and hands and feet. Not all cases will have symptoms.	Where rare additional symptoms develop e.g., high fever, headache, stiff neck, back pain or other complications, prompt medical advice should be sought.	Exclusion is not necessary, and cases can return to school as soon as they feel well.
Headlice	Other than the detection of live lice or nits, there are no immediate symptoms	Treatment is only necessary when live lice are seen.	Exclusion is not necessary, as headlice are not considered a health hazard.

Disease	Symptoms	Considerations	Exclusion Period
	until two to three weeks after infection, where itching and scratching of the scalp occurs.	Staff are not permitted to inspect any pupil's hair for headlice. If a staff member incidentally notices headlice in a pupil's hair, they will inform the pupil's parents and advise them to treat their child's hair. Upon noticing, staff members are not required to send the pupil home; the pupil is permitted to stay in school for the remainder of the day. When a pupil has been identified as having a case of headlice, a letter will be sent home to all parents notifying them that a case of headlice has been reported and asking all parents to check their children's hair.	In severe, ongoing cases, the LA does have the power to exclude. This use of power must be carefully considered, and exclusion should not be overused.
Hepatitis A	Infection can be asymptomatic. Symptoms can include abdominal pain, loss of appetite, nausea, fever and fatigue, followed by jaundice, dark urine and pale faeces.	The illness in children usually lasts one or two weeks, but can last longer and be more severe in adults.	Cases are excluded while unwell and for seven days after the onset of jaundice (or the onset of symptoms if no jaundice present).
Hepatitis B	Infection can be asymptomatic. Symptoms can include general fatigue, nausea, vomiting, loss of appetite, fever and dark urine, and older cases may develop jaundice. It can cause an acute	The HPT will be contacted where advice is required. The procedures for dealing with blood and other bodily fluids will	Acute cases will be too ill to attend school and their doctor will advise when they are fit to return. Chronic cases will not be excluded or

Disease	Symptoms	Considerations	Exclusion Period
	or chronic illness.	always be followed. The accident book will always be completed with details of injuries or adverse events related to cases.	have their activities restricted. Staff with chronic hepatitis B infections will not be excluded.
Hepatitis C	Symptoms are often vague but may include loss of appetite, fatigue, nausea and abdominal pain. Less commonly, jaundice may occur.	The procedures for dealing with blood and other bodily fluids will always be followed. The accident book will always be completed with details of injuries or adverse events related to cases.	Cases will not be excluded or have their activities restricted.
Impetigo	Symptoms include sores, typically on the face and on the hands and feet. After around a week, the sores burst and leave golden brown crusts, and can sometimes be painful and itchy.	Towels, facecloths and eating utensils will not be shared by pupils. Toys and play equipment will be cleaned thoroughly; non-washable soft toys will be wiped or washed with a detergent using warm water, and dried thoroughly.	Cases will be excluded until all sores or sores or blister are crusted over, or 48 hours after commencing antibiotic treatment.
Influenza	Symptoms include headache, high temperature, cough, sore throat, aching muscles and joints and fatigue. Younger cases may present different symptoms e.g., without fever but with diarrhoea.	Those in risk groups will be encouraged to have the influenza vaccine. Anyone with flu-like symptoms will stay home until they have recovered. Pupils under 16 will not be given aspirin.	There is no specific exclusion period; cases will remain home until they have fully recovered.
Invasive Group A Streptococcus	Symptoms include flu-like symptoms, sore throat, rough rash, scabs and sores	These infections are caused by the bacteria getting into parts of the	Inform HPT if any cases reported. The HPT will carry out a risk assessment and

Disease	Symptoms	Considerations	Exclusion Period
(iGAS)	(impetigo), pain and swelling, severe muscle aches, nausea and vomiting.	body where it is not normally found, such as the lungs or bloodstream. In rare cases an iGAS infection can be fatal.	undertake appropriate investigations and/ or actions as required.
Measles	Symptoms include a runny nose, cough, conjunctivitis, high fever and small white spots inside the cheeks. Around the third day, a rash of flat red or brown blotches may appear on the face then spread around the body.	All pupils are encouraged to have MMR immunisations in line with the national schedule. Staff members should be up-to-date with their MMR vaccinations. Pregnant staff members and those with weak immune systems will be encouraged to contact their GP immediately for advice if they come into contact with measles.	Cases are excluded while infectious, which is from four days before the onset of a rash to four days after.
Meningitis	Symptoms include fever, severe headaches, photophobia (aversion to light), stiff neck, non-blanching rash, vomiting and drowsiness.	Pupils are encouraged to be up-to-date with their vaccinations. Meningitis is a notifiable disease.	Once a case has received any necessary treatment, they can return to school once they have recovered.
Meningococcal Meningitis and Septicaemia (Sepsis)	Symptoms include fever, severe headache, photophobia, drowsiness and a non-blanching rash. Not all symptoms may be present.	Medical advice will be sought immediately. The confidentiality of the case will always be respected. The HPT and school health advisor will be notified of a case of meningococcal disease in the school. The HPT will be notified if two cases	When the case has been treated and recovered, they can return to school. Exclusion is not necessary for household or close contacts unless they have symptoms suggestive of meningococcal infection.

Disease	Symptoms	Considerations	Exclusion Period
		of meningococcal disease occur in the school within four weeks.	
Methicillin Resistant Staphylococcus Aureus (MRSA)	Symptoms are rare but include skin infections and boils.	All infected wounds will be covered.	No exclusion is required.
Mpox (monkeypox)	Symptoms are rare but begin within 5 to 21 days after a close physical contact with someone who has mpox infection and may include flu-like symptoms, fever, low energy, swollen glands and general body aches.	The case will be encouraged to consult their GP. Any close contacts will be advised to consult with their local HPT for advice.	Cases will be excluded until the rash has scabbed, all the scabs have fallen off and a fresh layer of skin has formed underneath.
Mumps	Symptoms include a raised temperature, swelling and tenderness of salivary glands, headaches, joint pain and general malaise. Mumps may also cause swelling of the testicles and ovaries.	The case will be encouraged to consult their GP. Parents are encouraged to immunise their children against mumps.	Cases can return to school five days after the onset of swelling if they feel able to do so.
Norovirus	Symptoms include nausea, diarrhoea and vomiting. It is known as the 'winter vomiting bug' and is the most common cause of gastroenteritis.	The HPT will be contacted if there are a higher than previously experienced and/ or rapidly increasing number of pupil and staff absences due to diarrhoea and vomiting.	Cases will be excluded for 48 hours after symptoms have stopped and they are well enough to return.
Panton-Valentine Leukocidin Staphylococcus Aureas (PVL-SA)	Symptoms can include recurrent boils, skin abscesses and cellulitis.	The HPT will be contacted if there are two or more cases.	Exclusion is not necessary unless cases have a lesion or wound that cannot be covered. Cases should not visit gyms or swimming pools until wounds have healed.
Respiratory Infections, including Cornoavirus	Symptoms can be caused by several respiratory infections including the common cold, coronavirus (COVID-19),	Pupils with symptoms will be encouraged to cover their mouth and nose with a tissue when	Cases with mild Symptoms e.g., a runny nose and/ or sore throat, can continue to attend if they are otherwise well.

Disease	Symptoms	Considerations	Exclusion Period
	flu and respiratory syncytial virus (RSV). Symptoms can be wide-ranging, including a runny nose, high temperature, cough and sore throat, and loss or change in sense of smell or taste.	coughing and sneezing, and to wash their hands afterwards. The DfE helpline and/ or the local HPT will be contacted if an outbreak occurs or there is evidence of severe disease, e.g., hospital admission.	Cases who are unwell and have a high temperature should remain at home until they no longer have a high temperature. It is not recommended that children and young people are tested for coronavirus unless directed to by a health professional. Cases aged 18 years and under with a positive test result should stay at home for 3 days after the day they took their test.
Ringworm	Symptoms vary depending on the area of the body affected. The main symptom is a rash, which can be scaly, dry, swollen or itchy and may appear red or darker than surrounding skin.	Pupils with ringworm of the feet will wear socks and trainers at all times and cover their feet during PE. Parents will be advised to consult their GP for recommended treatment.	No exclusion is usually necessary. For infections of the skin and scalp, cases can return to school once they have started treatment.
Rotavirus	Symptoms include severe diarrhoea, stomach cramps, vomiting, dehydration and mild fever.	Cases will be sent home if unwell and encouraged to speak to their GP.	Cases will be excluded until 48 hours have passed since symptoms were present.
Rubella (German Measles)	Symptoms are usually mild. They can include a rash, swollen lymph glands, sore throat and a runny nose, mild fever, headache, tiredness, conjunctivitis and painful swollen joints.	MMR vaccines are promoted to all pupils.	Cases will be excluded for five days from the appearance of the rash.
Scabies	Symptoms include tiny pimples and nodules on the skin. Burrows may be present on the wrists, palms, elbows, genitalia and buttocks.	All household contacts and any other very close contacts should have one treatment at the same time as the second treatment of the infected individual.	Cases will be excluded until after the first treatment has been carried out.

Disease	Symptoms	Considerations	Exclusion Period
		The second treatment must not be missed, and should be carried out one week after the first treatment.	
Scarlet Fever and Invasive Group A Streptococcal Disease	Scarlet Fever is highly infectious. It is usually a mild illness, though severe complications can occur in rare circumstances. It may be confused with measles. Symptoms can include: – Flu-like symptoms e.g., fever, swollen glands and an aching body – Sore throat and/ or tonsillitis – A rash that feels rough, like sandpaper i.e., scarlet fever, typically on the chest and stomach – Flushed cheeks – Scabs and sores – A white coating on the tongue	Cases will be encouraged to visit their GP. The HPT will be contacted if: – Two or more cases occur within 10 days of each other, and the affected individuals have a link – There are cases of serious disease which have resulted in overnight stays in hospital – There are cases of chickenpox and/ or influenza co-circulating in the group where a case of scarlet fever has been confirmed	Cases are excluded and can return 24 hours after commencing appropriate antibiotic treatment – cases not receiving treatment will be excluded until resolution of symptoms.
Slapped Cheek Syndrome, Parvovirus B19, Fifth's Disease	Where symptoms develop, a rose-red rash making the cheeks appear bright red may appear several days after a mild feverish illness. The rash usually peaks after a week and then fades.	Cases will be encouraged to visit their GP. Parents are requested to inform the school of a slapped cheek syndrome diagnosis.	Exclusion is not required – cases are not infectious by the time the rash occurs.
Threadworm	Symptoms include itching around the anus and vagina, particularly at night, and worms may be seen in stools or around the bottom.	Cases will be encouraged to visit their pharmacy for advice on treatment.	Exclusion is not required.
Tuberculosis (TB)	Symptoms include cough, loss of appetite, weight loss, fever, sweating	Advice will be sought from the HPT before taking any action, and	Cases with infectious TB can return to school after two weeks of treatment if

Disease	Symptoms	Considerations	Exclusion Period
	(particularly at night), breathlessness and pains in the chest. TB In parts of the body other than the lungs may produce a painful lump or swelling.	regarding exclusion periods.	well enough to do so, and so long as they have responded to anti-TB therapy. Cases with non-pulmonary TB, and cases with pulmonary TB who have effectively completed two weeks of treatment as confirmed by TB nurses, will not be excluded.
Typhoid and Paratyphoid Fever	Symptoms include fatigue, fever and constipation. The symptoms of Parathyroid Fever include fever, diarrhoea and vomiting.	All cases will be immediately reported to the HPT.	Cases will be excluded whilst symptomatic and for 48 hours after symptoms have resolved. Environmental health officers of the HPT may advise the school to issue a lengthened exclusion period.
Whooping Cough (Pertussis)	Symptoms include a heavy cold with a temperature and persistent cough. The cough generally worsens and develops the characteristic 'whoop'. Coughing spasms may be worse at night and may be associated with vomiting.	Cases will be advised to see their GP. Parents will be advised to have their children immunised against whooping cough.	Cases will be excluded from school until they have received 48 hours of appropriate treatment with antibiotics and feel well enough to return to school, or 21 days from the onset of illness if no antibiotic treatment is given. Cases will be allowed to return in the above circumstances, even if they are still coughing.

Appendix 2 – Infection Absence Periods

This table details the minimum required period for staff and pupils to stay away from school following an infection, as recommended by the UK Health Security Agency.

* Identifies a notifiable disease. It is a statutory requirement that doctors report these diseases to their local PHE centre.

Infection	Recommended minimum period to stay away from school	Comments
Athlete's Foot	None.	Advise cases to visit their local pharmacy or GP for advice and treatment. Individuals should not be barefoot at their setting e.g., in changing areas, and should not share towels, socks or shoes with others.
Chicken Pox	At least 5 days from onset of rash and until all blisters have crusted over.	Pregnant staff contacts should consult with their GP or midwife.
Cold Sores	None.	Avoid contact with the sores.
Conjunctivitis	None.	Advise cases to visit their local pharmacy or GP for advice and treatment. If an outbreak occurs, consult the HPT.
Respiratory Infections including Coronavirus (COVID 19)	Cases should not attend if they have a high temperature and are unwell. Cases who have a positive test result for COVID-19 should not attend the setting for 3 days after the day of the test.	Cases with mild symptoms such as runny nose and headache, who are otherwise well, can continue to attend.
Diarrhoea and/ or vomiting	Whilst symptomatic and 48 hours from the last episode.	Contact the HPT if there are a higher than previously experienced and/ or rapidly increasing number of absences due to diarrhoea and vomiting. If a particular cause of the diarrhoea and vomiting is identified, there may be additional exclusion advice, for example E. coli STEC and Hep. A.
Diphtheria *	Exclusion is essential.	Family contacts must be excluded until cleared by the HPT and the HPOT must always be consulted.

Infection	Recommended minimum period to stay away from school	Comments
Flu (Influenza)	Until recovered.	Report outbreaks to the HPT.
Glandular Fever	Exclusion is not necessary, and cases can return to school as soon as the feel well enough to do so.	Glandular Fever is spread through spit, and can be transferred through kissing or by sharing cups or cutlery. Cases will be infectious for up to 7 weeks before symptoms appear.
Hand, Foot and Mouth Disease	None.	Contact the HPT if a large number of children are affected. Exclusion may be considered in some circumstances.
Headlice	None.	Treatment recommended only when live lice seen. Exclusion is not normally permitted. In severe, ongoing cases, the LA does have the power to exclude; however, exclusion should not be overused.
Hepatitis A*	Seven days after onset of jaundice or other symptoms.	If it is an outbreak, the HPT will advise on control measures.
Hepatitis B*, C* and HIV	None.	Not infectious through casual contact. Procedures for bodily fluid spills must always be followed.
Impetigo	48 hours after commencing antibiotic treatment, or when lesions are crusted and healed.	Antibiotic treatment is recommended to speed healing and reduce the infectious period.
Measles*	Four days from onset of rash and when feeling well enough.	Preventable by vaccination (MMR). Follow procedures for vulnerable children and pregnant staff.
Meningococcal Meningitis*/ Septicaemia*	Until recovered.	Meningitis ACWY and B are preventable by vaccination. The HPT will advise on any action needed.
Meningitis*- due to other bacteria	Until recovered.	Hib and pneumococcal meningitis are preventable by vaccination. The HPT will advise on any action needed.
Meningitis* - viral	None.	As this is a milder form of meningitis, there is no reason to exclude those who have been in close contact with infected persons. The local HPT should be consulted.
MRSA	None.	Good hygiene – in particular environmental cleaning and handwashing – are important to minimise the spread. The local HPT should be consulted.
Mumps*	Five days after onset of swelling.	Preventable by vaccination with two doses of MMR.
Ringworm	Exclusion is not usually required.	Treatment is required.

Infection	Recommended minimum period to stay away from school	Comments
Rubella* (German Measles)	Five days from onset of rash.	Preventable by two doses of immunisation (MMR). Pregnant staff contacts should seek prompt advice from their GP or midwife.
Scabies	Can return to school after first treatment.	The infected person's household and those who have been in close contact will also require treatment at the same time.
Scarlet Fever* and Invasive Group A Streptococcal Disease	24 hours after commencing antibiotic treatment.	Antibiotic treatment is recommended, as a person is infectious for two to three weeks if antibiotics are not administered. If two or more cases occur, the HPT should be contacted.
Slapped Cheek Syndrome, Parvovirus B19, Fifth's Disease	None (once rash has developed).	Pregnant contacts of case should consult with their GP or midwife.
Threadworms	None.	Treatment recommended for the infected person and household contacts.
Tonsillitis	None.	There are many causes, but most causes are virus-based and do not require antibiotics.
Tuberculosis (TB)	If pulmonary TB – until at least 2 weeks after the start of effective antibiotic treatment. Exclusion not required for non-pulmonary or latent TB infection.	Only pulmonary (lung) TB is infectious. It requires prolonged close contact to spread. Consult the local HPT before disseminating information to staff and pupil families. The HPT will organise any necessary contact tracing.
Warts and Verrucae	None.	Verrucae should be covered in swimming pools, gymnasiums and changing rooms.
Whooping Cough* (Pertussis)	Two days from commencing antibiotic treatment, or 21 days from the onset of illness if no antibiotic treatment is given.	Preventable by vaccination. Non-infectious coughing can continue for many weeks after treatment. The HPT will organise any necessary contact tracing.

Appendix 3 – Diarrhoea and Vomiting Outbreak Action Checklist

Date	
Completed By	

Action	Action Taken?		Comments
	Yes	No	
A 48 hour exclusion rule has been enforced for ill pupils and staff			
Individuals with symptoms have been kept in an area away from communal areas where they can be observed until parent collects them.			
Liquid soap and paper hand towels are available at all hand wash basins.			
Enhanced cleaning is undertaken twice daily as a minimum, and an appropriate disinfectant is used.			
Advice has been given on the cleaning of vomit, e.g., steam cleaning carpets and furniture and machine hot washing of soft furnishings.			
Appropriate disposable personal protective equipment (PPE) is available.			
Appropriate waste disposal systems are in place for removing infectious waste.			
Appropriate spill kit is in place. Staff wearing appropriate PPE when dealing with spills, which will be removed and disposed of quickly.			
Hard toys are cleaned and disinfected on a daily basis, and their use is limited and rotated.			
The use of soft toys, water and sand play, and cookery activities have been suspended.			
Infected linen is segregated, and dissolvable laundry bags are used where possible.			
Visitors are restricted, and essential visitors are informed of the outbreak and advised on hand washing.			
New pupils joining the affected class of year group are delayed from joining.			

The health protection team (HPT) have been informed of any infected food handlers.			
Staff work in dedicated areas and food handling is restricted where possible.			
All staff (including agency) are asked if they are unwell and excluded for 48 hours if unwell.			
Staff work in dedicated areas where possible.			
Trays of fruit/ snacks covered until point of serving. Snacks served in individual bowls handed directly to pupils.			
Drink bottles clearly labelled with names.			
The HPT is informed of any planned events at the school.			